

Preservation of Malolactic Starters of *Lactiplantibacillus plantarum* strains Obtained by Solid-State Fermentation on Apple Pomace

Marina Edith Navarro ¹, Natalia Soledad Brizuela ¹, Naiquén Elizabeth Flores ¹, Manuel Morales ², Liliana Carmen Semorile ¹, Danay Valdes La Hens ¹, Adriana Catalina Caballero ², Barbara Mercedes Bravo-Ferrada ¹ and Emma Elizabeth Tymczyszyn ^{1,*}

¹ Laboratorio de Microbiología Molecular, Instituto de Microbiología Básica y Aplicada (IMBA), Departamento de Ciencia y Tecnología, Universidad Nacional de Quilmes, Bernal, CP 1876 Argentina; marina.edith.navarro@gmail.com (M.E.N.); nsbrizuela@gmail.com (N.S.B.); naiflores9007@gmail.com (N.F.); lsemorile@unq.edu.ar (L.S.); dvhens@gmail.com (D.V.L.H.); barbaramercedes@hotmail.com (B.B.-F.)

² Instituto de Investigación y Desarrollo en Ingeniería de Procesos, Biotecnología y Energías Alternativas (PROBIEN), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Universidad Nacional del Comahue (UNCo); Neuquen CP Q8300 Argentina. mangelmorales201213@gmail.com (M.M.); adrianacaballerobis@gmail.com (A.C.)

* Correspondence: elitym@yahoo.com.ar; Tel.: +54-011-4356-7100 (ext. 5636)

Supplemented Material




Figure S1. Cell viability (left) and percentage of malic acid consumption (% MAC, right) in synthetic wine fermentations (pH 3.8 and 12% ethanol) for 12 days, inoculated with freeze-dried UNQLp11 (●) or UNQLp155 (▲). Cultures were grown in MRS, then acclimated with ethanol 6% v/v and pH 4.5 for 48 h, according Brizuela et al 2017. Trehalose was added as cryoprotective agent and dried cultures were rehydrated in MRS at 21 °C for 24 h previous to wine inoculation. The results of %MAC were fitted according to a non-linear regression of exponential decay (Equation 2).

Author Contributions: Formal analysis, M.E.N., B.B.-F., and E.E.T.; Investigation, M.E.N., N.S.B., M.M., N.F., D.V.L.H., and A.C.; Methodology, M.E.N., N.S.B., and B.B.-F.; Project administration, L.S. and E.E.T.; Writing—original draft, M.E.N., B.B.-F., and E.E.T.; Writing—review & editing, M.E.N., N.S.B., B.B.-F., D.V.L.H., A.C., L.S., and E.E.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by grants from Agencia Nacional de Promoción Científica y Técnica (ANPCyT; PICT 2020 N° 3372, PICT 2019 N° 0008, PICT 2021 ApliCatI 0013) and the Con-

sejo Nacional de Investigaciones Científicas y Técnicas (CONICET) 35
(PIP-2021–11220200100394CO), Argentina. 36

Data Availability Statement: The original contributions presented in the study are included in the 37
article/supplementary material, further inquiries can be directed to the corresponding author. 38

Acknowledgments: N.S.B., B.B.-F., E.E.T., and M.M. are members of the Research Career of the 39
Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). N.F. is a fellow of CIC–PBA. 40
L.S. and D.V.L.H. are members of the Research Career of the CIC-PBA. 41

Conflicts of Interest: The authors do not have any conflicts of interest. 42

References 43

1. Brizuela, N.S.; Bravo-Ferrada, B.M.; La Hens, D.V.; Hollmann, A.; Delfederico, L.; Caballero, A.; Tymczyszyn, E.E.; Semorile, L. Comparative vinification assays with selected Patagonian strains of *Oenococcus oeni* and *Lactobacillus plantarum*. *LWT* **2017**, *77*, 348–355. 44
45
46
47